

MEDICATIONS TO AVOID BEFORE SURGERY

The following drugs contain aspirin or other products which can cause increased bleeding during surgery and should not be taken for 10 days prior to surgery. For relief of minor pain, you MAY take Tylenol (acetaminophen).

Advil	Coricidin	4 Way Cold Tablets	Oxaprozin
Aggrenox	Coumadin	Goody's Headache	Percodan
Aleve	Dabigatran	(Powder or Tablets)	Persantine
Alka Seltzer	Darvon	Herbal Supplements	Piroxicam
Alcohol	Darvon with Aspirin	Ibuprofen	Plavix
Anacin	Daypro	Indomethacin	Pradaxa
Anaprox	Diclofenac	Indocin	Quagesic
Ansaid	Diflunisal	Ketoprofen	Relafen
Arthotec	Disalcid	Lodine	Robaxisal
Aspirin	Doan's (Regular & Extra	Lovenox	Rufen
APC	Strength)	Meclomen	Sine Aid
Apixaban	Dolobid	Medipren	Soma
BC Tablets or Powders	Dristan	Meloxicam	Sulindac
BC Cold Powder	Duradyne Tablets	Midol 200	Trandate
Brilinta	Easprin	Midol PMS Caplets	Trental
Brufen	Ecotrin	Mobic	Trilisate
Bufferin	Eliquis	Motrin	Vanquish
Cama Arthritis Pain	Empirin	Naprelan	Vitamin E
Cataflam	Enbrel	Nabumeton	Yoltaren
Celebrex	Equagesic Tablets	Naprosyn	Warfarin
Compound	Etodolac	Naprosyn Pepto Bismol	Wesprin
Cephalgesics	Excedrin	(Tablets & Liquid)	Xarelto
Cilostazol	Feldene	Naproxen	Zavtrin
Clinoril	Fiorinal	Norgesic Forte	Zoprin
Cogesprin	Fish Oil	Omega-3	
	Flurbiprofen Sodium	Orudis	
		Oruvail	

Before stopping any medications, be sure to consult the physician who ordered them. Some, such as Coumadin (Warfarin) and Plavix, are ordered to prevent or treat serious conditions such as deep venous thrombosis, pulmonary embolisms, and other heart problems. **This is not an all-inclusive list.** If you are unsure if you are taking an aspirin product or an anti-inflammatory, please ask your doctor, nurse or pharmacist.